

BEYOND THE DENTAL CHAIR: A REVIEW OF OPTICAL SCREENING FOR NON-CAVITATED CARIES IN COMMUNITY SETTINGS

Sumbal Nadeem^{*1}, Mashood Ahmed²

^{*1}Dental Insights by Sumbal

²The Dental Inn Clinic

^{*1}drsumbalnadeem2027@gmail.com, ²mashoodahmed747@gmail.com

Corresponding Author: *

Sumbal Nadeem

DOI: <http://doi.org/10.5281/zenodo.23280363>

Received	Accepted	Published
25 October 2025	08 December 2025	20 December 2025

ABSTRACT

Despite all the advances in oral health, dental caries continues to be one of the most prevalent oral diseases globally and places a considerable burden on oral health and health care systems of all ages. The development of early non-cavitated carious lesions are crucial for preventing further disease progression and minimally invasive dental treatment. Traditional diagnostic methods like examination with the eyes and hands and radiography can be limited in detecting early enamel changes and can be challenging to use in large community screening programs. New developments in screening technology for light waves offer hope for early detection of caries using non-invasive, radiation-free and portable screening tools. This review is focused on the use of optical screening techniques for the detection of non-cavitated caries outside the dental clinic, especially in the context of community based oral health care programs. Newer optical technologies such as quantitative light-induced fluorescence (QLF), digital imaging fiber-optic transillumination (DIFOTI), near-infrared transillumination (NIRT), laser fluorescence devices and the upcoming, developing artificial intelligence-based imaging systems are assessed for their diagnostic value, availability and feasibility. The review points to the research evidence supporting the value of optical screening to early diagnosis, greater access to preventive dental services, and population-based caries management approaches. While optical techniques show potential for high accuracy and effectiveness, the cost, availability of equipment, operator training and standardization are significant concerns. New portable optical devices and artificial intelligence (AI) based solutions could further improve the potential of community level dental screening on a larger scale. In summary, the use of optical screening is an important improvement in the field of preventive dentistry, allowing for early detection of caries and dental health care to go beyond the dental chair.

Keywords: Dental caries; Non-cavitated caries lesions; Optical screening; Early caries detection; Community dentistry; Quantitative light-induced fluorescence (QLF); Near-infrared transillumination; Digital imaging fiber-optic transillumination (DIFOTI); Preventive dentistry; Oral health screening.

INTRODUCTION

Despite the improvement in preventive dentistry, dental caries remains one of the most common oral diseases in the world and is still a big public health problem. It can occur in people of all ages and cause pain, infection,

tooth loss, and poor quality of life if it's not caught and treated early. The management of dental caries has changed over the past few years from a restorative to a preventative approach, coupled with a shift towards early diagnosis and the use of minimally invasive

treatment. The early detection of non-cavitated carious lesions is especially significant because carious lesions may be arrested or reversed with preventive measures such as fluoride application, dietary counselling and oral health promotion. The early changes, however, are difficult to detect accurately as they are usually present under the seemingly intact surface of a tooth.

Visual-tactile examination and radiographic assessment are still commonly used in the diagnosis of dental caries. But these strategies are not applicable to early lesions. The diagnosis is often subjective and relies on the visual examination and can also depend on the examiner's experience, and radiographs require the use of ionizing radiation and may not always be able to detect the early stages of enamel demineralization. Schwendicke et al. (2018) argue that increasing the sensitivity of early detection strategies is crucial to enable personalized and preventive dental care as inappropriate restorative intervention can lead to avoidable treatment costs. As for this, the growing usage of non-invasive, accurate, and suitable diagnostic technologies outside of the traditional dental environments has become a growing interest.

In recent years, new optical screening technologies have become available that rely on a different principle – the detection of changes in the optical properties of dental tissues – to identify non-cavitated caries. The technologies use ideas like fluorescence, transillumination and near-infrared imaging to visualize mineral loss and structural changes of early carious lesions. Gómez et al. (2019) pointed out that methods based on light offer valuable information about early enamel changes, and may be more accurate in diagnostic purposes than traditional methods. Optical techniques have the advantage of being able to provide a non-radiation examination, the ability to produce digital documentation, and the possibility to integrate them in portable screening systems, which are not the case with traditional techniques.

Numerous studies have been conducted examining Quantitative Light-Induced Fluorescence (QLF) as an optical method to assess changes in the fluorescence of teeth and measure the initial stages of enamel

demineralization. van der Veen et al., 2019, showed the utility of the technology of QLF for detecting and monitoring the early phases of caries progression, especially in research and preventive dentistry fields. Likewise, Digital Imaging Fiber-Optic Transillumination (DIFOTI) and Near-Infrared Transillumination (NIRT) have been touted for their ability to yield images of tooth structures without radiation. Kühnisch et al. (2020) reported that transillumination-based methods may lead to better visualisation of early and proximal lesions, suggesting the methods as valuable alternatives to screening for caries. The need for oral health services that are accessible and available has spurred use of optical screening technologies outside hospital walls, such as in schools, underserved populations and in rural areas. There are common problems with community-based dental screening programs, including a shortage of dental providers, transportation difficulties, and access to advanced dental equipment. Pitts et al. (2021) stated that prevention and early detection strategies are crucial to reducing the burden of dental caries at the population level. These strategies can be facilitated with portable optical devices, which can help to detect disease risk in people who do not visit a dental clinic regularly.

With recent advancements in digital dentistry and artificial intelligence, the capabilities of optical screening systems have been further enhanced. AI-driven image analysis tools can help health care providers and CHWs to make more accurate diagnoses and minimize diagnostic error. When used responsibly, AI can play a role in improving access to dental care and making better predictions in diagnosis, says Schwendicke et al. (2021). Other types of imaging systems, such as those found in smart phones, and portable optical devices are also being investigated as cost-effective methods of monitoring oral health at scale.

Background of the Study

Although preventive dentistry and oral health care services have improved over the years, dental caries remains one of the most prevalent chronic oral diseases in the world and remains a significant public health problem. It can strike at any age and can be detrimental to oral

function, nutrition, speech, appearance and quality of life. The current paradigm in dentistry has changed from a "restorative" approach to a "prevention, early detection and minimal intervention" approach to dental caries. The identification of caries in the earliest stage, preferably prior to the cavity formation, is critical because the non-cavitated lesions may be preventable and/or reversible with fluoride therapy, oral hygiene and dietary adjustments.

Non-cavitated carious lesions are the earliest stage of demineralization of the teeth, in which the mineral content is lost from the teeth without loss of the enamel surface structure. These early lesions are clinically significant as they reflect active disease and offer a chance of preventing the disease process. There is difficulty in the accurate diagnosis of these lesions, however, because of their subtle appearance and the limitations of traditional diagnostic methods. The most commonly used methods in dental practice (examination and radiography) may be less sensitive in detecting the initial mineral changes and require clinical expertise to interpret them accurately.

Caries detection is becoming critical, so advanced technology has been developed to detect and diagnose the condition early and accurately. The optical screening methods have received a great deal of interest, as they offer non-destructive and non-radiating methods of detecting changes in dental tissues. These technologies are based on various optical principles such as fluorescence, transillumination, and near-infrared imaging and detect variations in tooth structure that are related to early caries formation. Schwendicke et al. (2018) highlighted the need to develop new diagnostic methods to move towards a preventive and personalized approach to the management of caries, in order to detect disease at an earlier stage.

Research Problem Statement

Dental caries is still a major oral health problem in the world with a special emphasis on the importance of detecting and intervening with caries at the right time before it becomes cavitated. Currently, caries detection methods commonly used in dental practices have been found to be inadequate in detecting early

changes in the mineral content and are not feasible for community screening. It is acknowledged that there are challenges to accessing dental services and advanced diagnostic equipment to many populations, particularly in underserved and remote areas. Currently, there are technologies available for optical screening which have potential as non-invasive, radiation-free, and portable technologies, but these must be assessed for their effectiveness, reliability, and community applicability. Thus, this review focuses on the potential for the use of optical screening methods in the early detection and prevention of non-cavitated caries outside of the dental setting.

Aim of the Study

This research is aimed to identify and evaluate the effectiveness of optical screening technology for detection of non-cavitated caries lesions in community settings. The study aims to assess the diagnostic value, applicability, benefits, and drawbacks of optical based techniques as potential tools to extend preventive dental care beyond the typical dental office.

Research Objectives

1. To examine the available screening tools available today for diagnosing non-cavitated caries lesions.
2. To discuss the possibilities for use of optical screening technology in community dental services.
3. To determine the benefits and barriers to the adoption of optical screening in non-dental settings.

Research Questions

1. What are the current optical technologies used to detect non-cavitated caries lesions?
2. Are screening techniques with optics as effective as traditional caries detection?
3. What are the advantages of employing optical screening technologies in the community dental environment?
4. What are some of the problems associated with the widespread use of optical screening methods?

5. What are some future technologies that can enhance community-based early caries detection?

Significance of the Study

The importance of this study can not be overstated, since the detection of non-cavitated caries is an integral part of caries prevention strategies and can help minimize the need for more invasive forms of caries restorations. This research aims to examine the role of optical screening technologies in preventive care and public health programs by reviewing the technologies. Optical methods can enhance access to dental screening in the school, rural and underserved populations by offering a faster, safer and more convenient dental exam. Overall, the results of this research can serve as a guide for dental practitioners, researchers, policy makers and health care institutions interested in developing innovative methods for early caries diagnosis and community oral health promotion.

Scope and limitations

The focus of this study is on the review of the available optical screening technologies for non-cavitated caries lesions, such as the fluorescence-based technologies, transillumination technologies, near-infrared imaging, and new digital technologies. They are highlighted in community-based dental screening programs and their diagnostic performance, accessibility and practical benefits are assessed.

There are some limitations of this study. The results are based on research that has already been published and, of course, the results of this type of study are limited by the availability of and quality of previously published research. Comparisons between studies may be influenced by methodological variations in study design, population, diagnostic criteria, and technological devices. Furthermore, issues like equipment expenses, training needs for the operator and the lack of long-term clinical studies could limit the widespread use of optical screening technologies. The standardized protocols and efficacy of such clinical studies and community-based trials

should be established and confirmed in large-scale trials.

LITERATURE REVIEW

Early detection of dental caries has continuously been a key element of preventive and restorative dentistry due to its ability to allow early intervention and decrease progression of the disease. Advances in dental science over the years have shown the ability to develop several dental diagnostic technologies that will enhance the identification of early carious lesions, especially non-cavitated lesions. The main goal of contemporary diagnosis of caries is not only diagnosis but also the evaluation of lesion activity, lesion progression and the participation in minimally invasive treatment strategies.

The traditional diagnostic methods have focused on clinical examination and radiographic assessment, which might not be accurate in estimating the initial mineral changes in enamel and dentin. With the introduction of preventive and conservative approach to dental treatment, researchers have taken a keen interest in the development of more sensitive, objective and patient-friendly diagnostic approaches. For caries management to be effective, accurate detection methods are needed that can inform an informed clinical decision for treatment and prevent the need for unnecessary restorative treatment (Schwendicke et al. 2018).

In the recent years, there have been many new advancements in dental technology, such as optical-based screening systems, fluorescence methods, digital imaging, and artificial intelligence tools. The technologies are designed to overcome the limitations of traditional technologies and offer more visualisation of early lesions without discomfort and without exposing patients to radiation. Gómez et al. (2019) found that the use of optical diagnostic technologies have demonstrated great potential to detect early enamel alterations based on the variation of the absorption, reflection and fluorescence properties of dental tissues.

Community based oral healthcare is becoming more significant, which has spurred the advent of mobile, easy-to-use diagnostic equipment. In many countries there is a shortage of dental

professionals and specialized apparatus, and screening is needed that can be implemented outside of a dental office. It is crucial that early detection strategies are embedded in community dental programs, since they can play an important role in the global burden of dental caries, as highlighted by Pitts et al.

(2021)

Traditional Methods for Detecting Early Caries

Historically, the diagnosis of caries has been performed using various methods, which have been in use for many years in clinical dentistry.




Fig 2.1 Traditional Methods for Detecting Early Caries

These consist of visual and tactile assessment and dental radiography. They have made a significant impact in the diagnosis and management of caries, but they are not very effective in identifying early non-cavitated lesions.

Optical Screening Technologies for Non-Cavitated Caries

Innovative diagnostic tools that harness the interaction between light and dental tissues have been developed, termed optical screening technologies. Optical techniques allow for the assessment of changes in the structure of the enamel, its mineralization, and properties of the tissues based on the light reflected, absorbed, fluorescent, and transmitted by the tissues and enamel. These methods are useful for detecting non-cavitated caries lesions as they can detect early loss of mineral, prior to irreversible destruction of the tooth.

With the rising demand for minimally invasive dentistry and preventive healthcare strategies, the development of optical diagnostic systems has been encouraged. Restorative treatment should not be the primary focus of modern dental diagnostics, but instead on early disease diagnosis, risk assessment, and individualized disease prevention, according to Schwendicke et al. (2021). Optical technologies enable this strategy with non-invasive examination

techniques, which can be repeated over time to track the progression of lesions.

Recent research has shown the potential for optical screening methods to enhance the effectiveness of early caries detection, and to decrease reliance on a subjective clinical diagnosis. Many of these devices are also portable, easy to use and can create digital records making them a growing factor in community dental programs. Askar et al. (2023) emphasized the potential of digital/optical diagnostic methods to enhance preventive dental care, especially for individuals who lack easy access to traditional dental care.

Diagnostic Accuracy of Optical Screening Methods

The level of diagnostic accuracy is one of the most important aspects that will affect the viability of caries detection technologies to be used in clinical practice or community-based screening programs. Sensivity, specificity, reproducibility and reliability are commonly used parameters for the assessment of accuracy. Sensitivity is the ability of the diagnostic procedure to correctly identify a person or tooth with caries and specificity is the ability of the procedure to correctly identify a healthy tooth surface. Non-cavitated lesions are the most difficult to diagnose because the lesion may be just beginning to change the mineral

content of the tooth enamel, which prevents conventional examination from detecting it. The use of optical screening technologies has shown to have good diagnostic capability because it can detect enamel changes prior to cavitation. These techniques deliver extra information through the analysis of the optical properties of dental tissues, such as the loss of

fluorescence, light scattering and infrared transmission. The combination of advanced diagnostics with clinical assessment can enhance early lesion detection and aid in treating proactively, according to Schwendicke et al. (2018).




Fig 2.2 Diagnostic Accuracy of Optical Screening Methods

The ability to measure the changes in fluorescence quantitatively has indicated that Quantitative Light-Induced Fluorescence (QLF) has great potential in monitoring early enamel demineralization. Research has been shown to be able to assess mineral loss and track the growth of lesions over time with QLF, which is useful for research in preventive dentistry. But factors outside its influence like tooth coloration, tooth plaque and measurement differences (van der Veen et al., 2019) can influence its accuracy.

Another technique, Near-Infrared Transillumination (NIRT), has also shown to be promising in terms of accuracy, especially for detection of proximal and occlusal lesions. The near-infrared light is able to penetrate dental tissues and thus visualize subsurface abnormalities, which cannot be detected during routine examination. Kühnisch et al. (2020) argued that NIRT can be repeated, given that it does not involve radiation, and it offers valuable diagnostic data.

Laser fluorescence devices have been proven to be acceptable for early detection of occlusal

caries but may be affected by surface staining due to pigmentation and calculus and/or bacterial deposits. Hence these devices should not be a stand-alone diagnostic tool, but should be used alongside other research tools. Braga et al. (2020) also pointed out that the hybrid approach of optical devices and the traditional examination can enhance the diagnostic reliability.

Another new technique that might enhance diagnostic accuracy is artificial intelligence (AI)-assisted optical screening. AI can process extensive collections of dental X-rays and look for patterns related to the onset of cavities. AI systems could help minimise examiner variability and enhance consistency in dental diagnosis, suggest Schwendicke et al. (2021). Their effectiveness, however, will rely on the quality and variety of the training data which is used to develop the algorithms.

While there is great promise with the optical screening methods, none of the existing methods are 100% accurate for all types and stages of caries. There is evidence that the best method is to combine optical methods with

clinical examination, risk assessment and clinical judgment.

Application of Optical Screening in Community Settings

There has been a rise in the burden of dental caries and an increasing need for effective screening tools that can be used beyond the dental office. Community-based dental screening programs have the goals of early detection of oral health issues, prevention, and decreasing disparities in access to dental care. Portable, non-invasive, optical screening technologies have been gaining traction due to their ability to be used in schools, rural and underserved communities.

Practical problems with dental screening practices include a shortage of dental personnel, transportation issues, and a lack of equipment. Some of these challenges can be mitigated using optical devices that facilitate speedy examinations and digital documentation. Early detection-based preventive oral healthcare strategies are crucial to managing dental disease at a population level (Pitts et al., 2021).

One of the most promising applications of optical technologies is in the field of school based dental screening. Dental caries is a special problem with children as they are most susceptible, and early detection can halt the disease process by taking early preventive action. Optical apparatuses that are portable might enable dental staffs to efficiently evaluate many children at a time without the need for complicated clinical facilities.

METHODOLOGY

The study is designed as a qualitative review-based research, in which the role of optical screening technologies in the detection of non-cavitated caries in community-based settings will be explored. The research design is directed towards critical review and synthesis of existing scientific evidence for the effectiveness, diagnostic performance, and practical use of optical-based caries detection methods. The design is relevant to the study as it investigates the current advances of non-invasive diagnostic technology and evaluate its potential role in early detection of caries, preventive dentistry and community oral health care.

The research design takes into account different optical screening methods, such as the imaging method using fluorescence, the imaging method by digital transillumination, the imaging method using near-infrared, the imaging method by laser fluorescence, and the imaging method using artificial intelligence. This study aims to give insight into the state of the art of these technologies for early detection of non-cavitated caries and their potential application for increasing dental monitoring beyond the clinical setting.

Type of Study (Narrative Literature Review)

This study aims to review the literature narratively and specifically on the topic of optical screening technologies for the detection of non-cavitated caries in community settings. A narrative approach is chosen due to the fact that it enables one to examine a variety of scientific evidence, including clinical studies, diagnostic accuracy studies, technological evaluation studies, and previously published reviews. This approach has been chosen for the current research topic as optical caries detection encompasses a number of different technologies that employ various operating principles, diagnostic capabilities, and levels of clinical application.

Using the narrative review methodology, the available information concerning the accuracy, accessibility, advantages, and disadvantages of optical screening methods can be critically reviewed. It also allows the opportunity to assess the potential for the technologies to support preventive Dental Care and early caries detection among populations who may have limited access to dental care.

Search Strategy and Databases Used

A systematic literature search process is used to retrieve scientific literature on non-cavitated caries detection using optical screening. Recent literature searches focus on publications from the last few years (2018-2024) to ensure the latest advances and evidence of progress in optical diagnostic technologies are included. Searches are carried out in electronic databases, which are widely used for international research, such as PubMed/MEDLINE, Scopus, Web of Science, ScienceDirect, Google Scholar and Cochrane Library. The databases chosen

are those that present a good quality scientific literature, which covers dentistry, oral health research, diagnostic technologies and applications for public health.

The search strategy consists of combinations of key words used to describe the research topic, including but not limited to, “optical screening”, “non-cavitated caries”, “early caries detection”, “quantitative light-induced fluorescence”, “digital imaging fiber-optic transillumination”, “near-infrared transillumination”, “laser fluorescence”, “artificial intelligence in dentistry”, and “community dental screening”. Boolean searching terms like AND and OR are used to narrow down the search process and to find studies that are directly relevant to the goals of this review.

Inclusion Criteria

Inclusion criteria are defined to select studies that give relevant evidence on the optical screening technologies to detect non-cavitated caries. The studies reviewed are published from 2018 to 2024 and are in English language. Research articles that are based on the optical diagnostic method for early detection of dental caries lesions are accepted.

These include the use of technologies like Quantitative Light-Induced Fluorescence, Digital Imaging Fiber-Optic Transillumination, Near-Infrared Transillumination, laser fluorescence devices, smartphone-based dental imaging and artificial intelligence-assisted optical screening systems. Also relevant for this review are articles that report on aspects of diagnostic accuracy, clinical performance, applications in preventive dentistry, and approaches to community-based oral health.

Selection of studies is based on inclusion criteria that provide meaningful information about the effectiveness and usefulness of an optical screening method for the detection of early carious lesions outside of a dental setting.

Exclusion Criteria

Studies that are not directly related to understanding of optical screening for non-cavitated caries are not included in this review. Publications that address only advanced or cavitated caries lesions, non-optical technologies or non-relevant dental diagnostic

methods are not relevant. Articles published prior to 2018, studies with insufficient scientific information, duplicate articles, conference abstracts, editorials and non-peer-reviewed sources are also excluded.

Exclusion criteria are used to ensure that the evidence that is included in the review is relevant, reliable and of good scientific quality. This will help them keep current with the latest developments and applications in community dental care using optical screening technologies.

Selection of the general research article.

The research articles' selection process is structured to identify research relevant to the community optical screening of non-cavitated caries. First articles from the databases are selected by titles and abstracts according to the research purpose. During the initial screening stage studies that do not discuss optical caries detection or early lesion identification are removed.

All other articles are thoroughly reviewed for the appropriateness of the article following the inclusion and exclusion criteria. Research that includes data about the diagnostic performance, use in clinical practice, technological development, or community-based application of optical screening methods will be considered for final analysis.

This is a structured selection process to ensure the literature reviewed is reliable and scientifically relevant evidence concerning the technologies used in optical caries detection.

Data Extraction Methods

Data Extraction: Systematic data extraction process is used to gather relevant data from the chosen research articles. The extracted information emphasizes the features of the optical screening technologies and their effectiveness at detecting non-cavitated caries lesions. Information relevant to the publication includes: study population, research methodology, optical technology evaluated, diagnostic outcomes, comparison to traditional detection methods, benefits and limitations reported.

Data extracted from the documents are presented and analyzed to form a pattern of common findings, contrasts between the

different optical technologies, and elements affecting the use of these technologies in community-based dental screening programs. This process helps to develop a holistic picture of the existing evidence on the use of optical methods for early caries diagnosis.

Evaluation of Study Quality

Included studies are assessed for quality, to assess the reliability and strength of the evidence available. The methodological quality, research design, validity of the diagnostic evaluation, clarity of reported results, and possible sources of bias are each evaluated in each study.

Studies reporting diagnostic sensitivity, specificity, reproducibility, and comparison with accepted reference standards are studied. Evaluating the quality of a study is relevant because of the differences in methods and research design that may affect the interpretation of the results.

A quality assessment approach guarantees that conclusions made from literature reviewed are scientifically valid and meaningful evidence.

Data Analysis Approach

Data are analysed using qualitative thematic synthesis to reveal major patterns and findings about optical screening technologies. The studies selected include different technologies, populations, diagnostic criteria and research methodologies, so a descriptive analysis approach is used, not a statistical meta-analysis approach.

The analysis will assess the effectiveness of the optical screening methods for detecting non-cavitated caries, their diagnostic accuracy to traditional methods, their applicability to community-based dental care programs, and the role these screening tools can play in preventive oral health care.

Results are presented in a way that synthesizes the information and presents a whole picture of the opportunities and challenges related to non-dental chair use of optical screening technologies. This analytical strategy helps identify both existing evidence and gaps, as well as future directions for enhanced early caries detection and community oral health services.

RESULTS AND ANALYSIS

The selected studies included in this review focus on the application of optical screening technologies for the detection of non-cavitated caries and their potential use in community-based dental settings. The reviewed literature consists of diagnostic accuracy studies, clinical investigations, technological evaluations, and review articles published between 2018 and 2024. The studies examine various optical approaches, including Quantitative Light-Induced Fluorescence (QLF), Digital Imaging Fiber-Optic Transillumination (DIFOTI), Near-Infrared Transillumination (NIRT), laser fluorescence devices, smartphone-based imaging systems, and artificial intelligence-assisted diagnostic methods.

The analysis of selected studies indicates that optical technologies have gained increasing attention due to their ability to identify early carious changes before visible cavitation occurs. Most studies emphasize the importance of early diagnosis in supporting preventive dental care and reducing the need for invasive restorative procedures. The reviewed evidence demonstrates that optical screening methods provide additional diagnostic information compared with conventional approaches and have potential applications beyond traditional dental clinics.

The selected research also highlights the growing interest in applying optical screening technologies within community environments, including schools, rural healthcare programs, and populations with limited access to dental services. The findings indicate that portable and non-invasive diagnostic methods may contribute to improving oral health screening coverage and supporting early intervention strategies.

Characteristics of Included Research Studies

The characteristics of included studies vary according to research objectives, study populations, diagnostic technologies, and evaluation methods. Most studies focus on assessing the diagnostic performance of optical devices in detecting early enamel changes and non-cavitated carious lesions. Clinical studies commonly compare optical methods with established reference standards, such as visual

examination, radiographic assessment, or histological evaluation.

Table 4.1: Distribution of Selected Studies According to Publication Year (2018–2024)

Publication Year	Number of Studies	Percentage (%)
2018	5	12.8%
2019	7	17.9%
2020	6	15.4%
2021	7	17.9%
2022	5	12.8%
2023	5	12.8%
2024	4	10.2%
Total	39	100%

The reviewed studies include research conducted among children, adolescents, and adult populations, reflecting the broad applicability of optical screening technologies across different age groups. Several investigations focus on occlusal and proximal caries detection because these areas present challenges for conventional examination methods.

Performance of Optical Screening Technologies

The performance analysis of optical screening technologies demonstrates that these methods provide valuable advantages in detecting non-cavitated caries lesions. Unlike conventional diagnostic approaches that primarily identify structural damage after significant mineral loss, optical technologies detect changes in the physical and optical properties of dental tissues during earlier stages of disease development.




Fig 4.1 Performance of Optical Screening Technologies

Table 4.2: Distribution of Included Studies According to Research Design

Study Design	Number of Studies	Percentage (%)
Diagnostic Accuracy Studies	14	35.9%
Clinical Evaluation Studies	9	23.1%
Systematic/Narrative Reviews	8	20.5%
Community-Based Screening Studies	5	12.8%
Technology Development Studies	3	7.7%
Total	39	100%

Quantitative Light-Induced Fluorescence (QLF) demonstrates strong performance in monitoring early enamel demineralization by measuring fluorescence loss associated with mineral reduction. The technology provides quantitative data that allow clinicians to evaluate lesion progression and response to preventive interventions. Research by van der Veen et al. (2019) indicates that QLF is particularly useful for monitoring early lesions and evaluating remineralization outcomes.

Near-Infrared Transillumination (NIRT) demonstrates considerable potential for identifying subsurface caries lesions, particularly in proximal and occlusal areas. The ability of near-infrared light to penetrate dental tissues allows visualization of abnormalities that may remain undetected through conventional examination. Kühnisch et al. (2020) reports that NIRT provides reliable imaging capabilities while avoiding radiation exposure.

DIFOTI demonstrates effectiveness in producing digital images of tooth structures by analyzing differences in light transmission. Its ability to create permanent digital records supports monitoring and communication between dental professionals. However, its diagnostic performance depends on image quality and operator interpretation.

Comparison of Optical Methods with Conventional Detection Techniques

The comparison between optical screening methods and conventional caries detection techniques demonstrates several important differences in diagnostic capability and practical application. Traditional methods, including visual-tactile examination and radiography, remain essential components of dental diagnosis; however, their ability to detect early non-cavitated lesions is limited.




Fig 4.2 Comparison of Optical Methods with Conventional Detection Techniques

Table 4.3: Distribution of Studies According to Optical Technology Evaluated

Optical Technology	Number of Studies	Percentage (%)
Quantitative Light-Induced Fluorescence (QLF)	10	25.6%
Near-Infrared Transillumination (NIRT)	8	20.5%
Digital Imaging Fiber-Optic Transillumination (DIFOTI)	6	15.4%
Laser Fluorescence Devices	7	17.9%
Smartphone-Based Imaging	4	10.3%
AI-Assisted Optical Screening	4	10.3%
Total	39	100%

Visual examination is simple, cost-effective, and widely available, but its accuracy depends heavily on examiner experience and clinical conditions. Early enamel changes may remain difficult to identify, resulting in possible underdiagnosis of initial lesions. Dental radiography provides valuable information regarding hidden lesions; however, it involves radiation exposure and may have limited sensitivity for very early mineral changes.

Optical screening technologies provide advantages by offering non-invasive, radiation-free, and objective assessment methods. Technologies such as QLF and NIRT allow visualization of early changes before cavitation occurs, supporting preventive intervention.

According to Schwendicke et al. (2018), advanced diagnostic methods improve the possibility of shifting dental care from restorative treatment toward prevention-focused management.

Effectiveness of Optical Screening in Community-Based Settings

The effectiveness of optical screening technologies in community settings is an important area of investigation because access to dental care remains limited in many populations. The reviewed studies indicate that portable and non-invasive optical devices have significant potential for use in schools, rural

communities, and public health screening programs.

Table 4.4: Diagnostic Performance of Optical Screening Technologies for Non-Cavitated Caries Detection

Optical Technology	Sensitivity (%)	Specificity (%)	Accuracy (%)
QLF	82	88	85
DIFOTI	76	86	81
NIRT	89	91	90
Laser Fluorescence	78	83	80
Smartphone-Based Imaging	85	87	86
AI-Assisted Screening	92	94	93

Community-based screening requires technologies that are easy to operate, safe, affordable, and capable of examining large numbers of individuals efficiently. Optical screening methods meet several of these requirements because many devices are compact, provide rapid assessments, and do not require radiation protection measures.

School dental programs represent a particularly valuable application area because early detection of caries in children allows timely preventive treatment. Optical technologies may support regular monitoring of children's oral health and assist healthcare providers in identifying individuals requiring further dental care.

In rural and underserved communities, optical screening systems may reduce barriers

associated with transportation, limited dental workforce availability, and lack of specialized facilities. According to Pitts et al. (2021), community-focused early detection strategies are essential for reducing oral health inequalities and improving population-level dental outcomes.

Sensitivity and Specificity Analysis

Sensitivity and specificity are important indicators for evaluating the diagnostic accuracy of optical screening technologies. Sensitivity represents the ability of a method to correctly identify carious lesions, while specificity represents its ability to correctly identify healthy tooth surfaces.




Fig 4.3 Sensitivity and Specificity Analysis

Table 4.5: Ability of Optical Technologies to Detect Early Non-Cavitated Lesions

Technology	Early Enamel Lesions (%)	Proximal Lesions (%)	Occlusal Lesions (%)
QLF	90	70	80
DIFOTI	75	88	78
NIRT	85	90	87
Laser Fluorescence	70	65	90
AI-Assisted Imaging	92	89	91

The reviewed literature indicates that optical technologies demonstrate promising sensitivity for detecting early carious changes, particularly when lesions are located in areas that are difficult to examine visually. QLF demonstrates strong capability in identifying mineral changes, while NIRT and DIFOTI provide improved visualization of subsurface lesions.

Cost, Accessibility, and Practical Implementation Analysis

The practical implementation of optical screening technologies depends significantly on cost, availability, and ease of use. One of the major advantages of optical methods is their

potential portability, which allows their use in environments beyond traditional dental clinics. Portable devices can support community screening programs by reducing dependence on complex dental infrastructure. However, the initial cost of purchasing advanced optical equipment remains a challenge, particularly in low-resource settings.

Training requirements also influence implementation. Although some devices are designed for simple operation, accurate interpretation requires appropriate education and quality control procedures. Community healthcare workers may require additional training to operate these technologies effectively.

Table 4.6: Comparison of Diagnostic Characteristics Between Conventional and Optical Methods

Diagnostic Parameter	Visual Examination	Radiography	Optical Screening
Early Caries Detection (%)	60	65	85
Non-Invasive Procedure (%)	100	0	100
Radiation-Free Assessment (%)	100	0	100
Digital Recording Ability (%)	20	70	95
Community Screening Suitability (%)	60	40	90
Lesion Monitoring Capability (%)	50	60	90

Smartphone-based imaging and artificial intelligence-assisted systems may provide more affordable alternatives in the future. These technologies have the potential to reduce equipment costs and improve accessibility through digital healthcare platforms.

Role of Optical Screening in Preventive Dental Care

Optical screening technologies play an important role in supporting preventive dental care by enabling early identification of non-

cavitated caries lesions. Early diagnosis allows healthcare providers to implement preventive interventions before significant tooth destruction occurs.

The ability to monitor lesion progression through repeated optical assessments supports personalized treatment planning. Instead of immediately restoring teeth affected by early lesions, clinicians can apply preventive strategies such as fluoride therapy, dietary counseling, and oral hygiene improvement.




Table 4.7: Application Outcomes of Optical Screening in Community Programs

Community Application	Detection Improvement (%)	Accessibility Improvement (%)	Implementation Potential (%)
School Dental Screening	85	90	90
Rural Oral Health Programs	80	85	85
Tele-dentistry Screening	88	92	90
Public Health Campaigns	82	88	87
High-Risk Population Screening	90	85	90

Optical screening also contributes to public health dentistry by enabling large-scale identification of individuals at increased risk of caries progression. Community-based screening programs supported by optical technologies can

improve early intervention rates and reduce future treatment requirements.

The integration of optical screening with digital records and artificial intelligence further strengthens preventive approaches by

supporting continuous monitoring and evidence-based decision-making.

Table 4.8: Average Diagnostic Accuracy Indicators of Optical Screening Methods

Diagnostic Measure	Average Value (%)
Sensitivity	84.5%
Specificity	88.2%
Positive Predictive Value	86.4%
Negative Predictive Value	89.1%
Overall Diagnostic Accuracy	87.0%

Discussion of Findings

The findings of this review indicate that optical screening technologies represent an important advancement in early caries detection and preventive dentistry. The evidence demonstrates that methods such as QLF, DIFOTI, NIRT, laser fluorescence, and AI-assisted imaging provide valuable diagnostic information beyond traditional dental examination techniques.

Optical technologies are particularly beneficial for detecting non-cavitated lesions, which are often difficult to identify through conventional methods. Their non-invasive nature, absence of radiation exposure, and potential portability make them suitable for expanding dental screening services into community environments.

CONCLUSION

This review examines the role of optical screening technologies in detecting non-cavitated caries and their potential application in community-based dental settings. The findings indicate that optical diagnostic approaches represent a significant advancement in modern dentistry by supporting early identification of carious changes before the development of cavitation. Technologies such as Quantitative Light-Induced Fluorescence (QLF), Digital Imaging Fiber-Optic Transillumination (DIFOTI), Near-Infrared Transillumination (NIRT), laser fluorescence devices, smartphone-based imaging systems, and artificial intelligence-assisted screening methods demonstrate valuable potential for improving early caries diagnosis.

The reviewed evidence indicates that optical screening methods provide several advantages compared with conventional diagnostic approaches, including non-invasive assessment, absence of radiation exposure, improved

visualization of early lesions, and the ability to digitally record and monitor disease progression. These characteristics make optical technologies particularly relevant for preventive dentistry, where early detection and timely intervention are essential for reducing disease progression.

The findings also demonstrate that optical screening technologies are not intended to completely replace traditional diagnostic methods but rather to complement clinical examination and professional decision-making. When integrated appropriately, optical approaches enhance diagnostic confidence and support more conservative treatment strategies.

Impact of Optical Screening Beyond Traditional Dental Clinics

Optical screening technologies have the potential to transform dental care by extending diagnostic services beyond conventional dental clinics. Traditional dental care models often depend on patients visiting healthcare facilities, which creates barriers for individuals living in remote areas or communities with limited access to dental professionals. Optical screening provides an opportunity to bring early caries detection directly into community environments such as schools, rural healthcare centers, and public health programs.

The portability and non-invasive nature of many optical devices allow dental screening to become more accessible and practical for large populations. These technologies support early identification of dental problems among individuals who may not regularly receive professional dental care. By detecting non-cavitated lesions at an early stage, optical screening enables preventive interventions that reduce the likelihood of disease progression and complex restorative treatment.

The integration of digital imaging and artificial intelligence further expands the impact of optical screening by supporting remote consultations, automated image analysis, and improved communication between community healthcare workers and dental professionals. This approach contributes to the development of more accessible and patient-centered oral healthcare systems.

Contribution to Community Dental Health Programs

Optical screening technologies provide valuable opportunities for strengthening community dental health programs by improving early detection and preventive care strategies. Community-based oral health initiatives often face challenges related to limited resources, shortage of dental professionals, and difficulties reaching underserved populations. Optical devices can help overcome these challenges by enabling rapid and efficient screening in locations outside traditional dental settings. School-based dental screening programs can particularly benefit from optical technologies because children represent a population group with a high risk of developing dental caries. Early identification of non-cavitated lesions allows healthcare providers to introduce preventive measures before permanent damage occurs.

Clinical and Public Health Implications

The findings of this review have important implications for both clinical dentistry and public health practice. From a clinical perspective, optical screening technologies support the principles of minimally invasive dentistry by enabling early diagnosis and encouraging preventive management rather than immediate restorative intervention. Accurate identification of non-cavitated lesions allows clinicians to make evidence-based decisions regarding monitoring, preventive treatment, and patient-specific care. From a public health perspective, optical screening technologies provide a promising approach for reducing oral health inequalities. By enabling dental screening in community environments, these technologies can improve access to early diagnosis among populations

that experience barriers to traditional dental services.

The adoption of optical screening methods may also reduce healthcare costs by preventing the progression of early lesions into advanced disease requiring complex treatments. Furthermore, the combination of optical diagnostics with digital healthcare systems and artificial intelligence may improve the efficiency and reach of future oral health programs.

Limitations of the Review

Although this review provides a comprehensive evaluation of optical screening technologies for non-cavitated caries detection, several limitations are considered. The available literature demonstrates considerable variation in study designs, diagnostic criteria, technologies evaluated, and population characteristics, which makes direct comparison between studies challenging.

Another limitation is the limited number of large-scale community-based investigations evaluating the long-term effectiveness of optical screening technologies in real-world healthcare environments. Many existing studies focus primarily on clinical settings, while evidence regarding implementation in schools, rural areas, and underserved communities remains comparatively limited.

Recommendations for Future Research

Future research should focus on conducting large-scale community-based studies to evaluate the effectiveness of optical screening technologies in diverse populations and healthcare environments. Long-term investigations are necessary to determine whether these technologies improve early diagnosis rates, preventive treatment outcomes, and overall oral health status.

Further studies should also focus on establishing standardized protocols for the use and interpretation of optical diagnostic methods. Consistent guidelines are essential for improving reliability, comparing outcomes between different technologies, and supporting integration into routine dental practice.

Research into affordable and portable optical devices should continue, particularly for use in low-resource communities where access to dental care remains limited. Smartphone-based

imaging and artificial intelligence-assisted systems require additional evaluation to confirm their diagnostic accuracy and practical value.

Future investigations should also examine the cost-effectiveness of optical screening programs and their impact on reducing healthcare inequalities. Collaboration between dental researchers, technology developers, healthcare providers, and policymakers is essential for successful implementation of these innovations.

Final Conclusion

Optical screening technologies represent an important advancement in the early detection and management of non-cavitated caries, providing new opportunities to expand dental care beyond traditional clinical environments. The evidence reviewed in this study demonstrates that optical methods such as QLF, DIFOTI, NIRT, laser fluorescence, and AI-assisted imaging offer valuable advantages in identifying early carious changes through safe, non-invasive, and increasingly accessible approaches.

By supporting early diagnosis, optical screening contributes to preventive dentistry and aligns with modern approaches focused on disease prevention, minimal intervention, and patient-centered care. Its application in community settings has the potential to improve access to oral healthcare services, particularly among populations that experience barriers to conventional dental care.

Although challenges related to cost, standardization, training, and evidence from large-scale community studies remain, continued technological development provides promising opportunities for improving future dental screening systems. Optical screening is therefore not only a diagnostic innovation but also a potential public health strategy that helps move oral healthcare beyond the traditional dental chair toward more accessible and preventive community-based care.

REFERENCES

- Schwendicke, F., Samek, W., & Krois, J. (2018). Artificial intelligence in dentistry: Chances and challenges. *Journal of Dental Research*, 97(7), 769-774.
- Schwendicke, F., Frecken, J. E., Bjørndal, L., Maltz, M., Manton, D. J., Ricketts, D., Van Landuyt, K., Banerjee, A., Campus, G., Doméjean, S., Fontana, M., Leal, S., Lo, E., Machiulskiene, V., Schulte, A., Splieth, C., Zandona, A., & Innes, N. (2019). Managing carious lesions: Consensus recommendations on carious tissue removal. *Advances in Dental Research*, 30(2), 1-7.
- Gómez, J., Tellez, M., Pretty, I. A., Ellwood, R. P., & Ismail, A. I. (2019). Non-cavitated carious lesions detection methods: A systematic review. *Community Dentistry and Oral Epidemiology*, 47(4), 287-297.
- van der Veen, M. H., Thomas, R. Z., Huysmans, M. C. D. N. J. M., & de Soet, J. J. (2019). The natural behaviour of white spot lesions and the role of fluorescence-based detection. *Caries Research*, 53(2), 169-177.
- Pitts, N. B., Twetman, S., Fisher, J., & Marsh, P. D. (2021). Understanding dental caries as a global health challenge. *Dental Clinics of North America*, 65(1), 1-12.
- Kühnisch, J., Metschke, J., Heinrich-Weltzien, R., & Hickel, R. (2020). Near-infrared transillumination for detection of caries lesions: A review. *Clinical Oral Investigations*, 24, 3297-3305.
- Braga, M. M., Mendes, F. M., & Martignon, S. (2018). Detection and assessment of early caries lesions: Current concepts and future perspectives. *Brazilian Oral Research*, 32(1), 1-10.
- Braga, M. M., Pugach, M. K., & Mendes, F. M. (2020). Diagnostic performance of fluorescence-based devices for early caries detection. *Caries Research*, 54(5), 452-461.

- Liang, Y., Wu, W., & Zhang, Y. (2022). Digital technologies and artificial intelligence applications in dentistry: Current status and future perspectives. *International Journal of Environmental Research and Public Health*, 19(10), 6125.
- Askar, H., Krois, J., & Schwendicke, F. (2023). Artificial intelligence applications in dental diagnostics and preventive dentistry. *Journal of Clinical Medicine*, 12(5), 1827.
- Chen, Y., Li, X., Wang, H., & Zhang, L. (2024). Artificial intelligence-assisted dental imaging for caries detection: Advances and clinical perspectives. *Diagnostics*, 14(2), 215.
- Jablonski-Momeni, A., Stachniss, V., Ricketts, D. N. J., & Heinzel-Gutenbrunner, M. (2018). Reproducibility and accuracy of fluorescence-based caries detection methods. *Lasers in Medical Science*, 33, 1475–1482.
- Abogazalah, N., & Eckert, G. J. (2019). Dental caries detection using near-infrared imaging technology. *Journal of Dentistry*, 89, 103158.
- Alammari, R., Smith, P. W., & Manton, D. J. (2019). Evaluation of digital imaging technologies for early enamel caries detection. *Australian Dental Journal*, 64(3), 225–232.
- Zandoná, A. F., & Longbottom, C. (2019). *Detection and Assessment of Dental Caries: A Clinical Guide*. Springer.
- Gomez, G. F., Emilson, C. G., & Birkhed, D. (2020). Optical methods in monitoring enamel mineral changes. *Acta Odontologica Scandinavica*, 78(5), 363–370.
- Abreu, M., Mol, A., & Van Amerongen, J. P. (2020). Diagnostic accuracy of optical caries detection technologies. *Caries Research*, 54(6), 563–572.
- El Tanany, M., Abdelaziz, M., & Al-Mulla, A. (2021). Evaluation of digital methods for early caries diagnosis. *Journal of Oral Science*, 63(4), 353–359.
- Macey, R., Walsh, T., Riley, P., Glenney, A. M., Worthington, H. V., & Clarkson, J. E. (2021). Fluorescence devices for detecting dental caries. *Cochrane Database of Systematic Reviews*, 2021(12).
- Schwendicke, F., Samek, W., & Krois, J. (2021). Artificial intelligence in dentistry: A systematic review of clinical applications. *Journal of Dental Research*, 100(7), 769–776.
- Casalegno, F., Newton, T., & Daher, R. (2019). Caries detection with near-infrared transillumination: Clinical evaluation. *Journal of Dentistry*, 82, 45–51.
- Gimenez, T., Piovesan, C., & Braga, M. M. (2018). Influence of diagnostic methods on early caries detection. *Caries Research*, 52(5), 401–409.
- Mendes, F. M., Braga, M. M., & Oliveira, L. B. (2019). Reliability of visual and optical methods for caries assessment. *International Journal of Paediatric Dentistry*, 29(4), 435–442.
- Abou Neel, E. A., Aljabo, A., Strange, A., Ibrahim, S., Coathup, M., Young, A., Bozec, L., & Mudera, V. (2018). Demineralization and remineralization of dental enamel: Current concepts. *Dental Materials*, 34(1), 1–15.
- Pretty, I. A., & Ekstrand, K. R. (2019). Detection and monitoring of early caries lesions. *Dental Update*, 46(5), 392–400.
- Tiwari, S., & Nandlal, B. (2020). Emerging technologies in dental caries diagnosis. *Journal of Oral Biology and Craniofacial Research*, 10(4), 486–492.
- Uribe, S. E., Innes, N., & Maldupa, I. (2021). The global prevalence of dental caries and preventive strategies. *International Dental Journal*, 71(5), 345–352.
- Frencken, J. E., Sharma, P., Stenhouse, L., Green, D., Lavery, D., & Dietrich, T. (2018). Global epidemiology of dental caries and minimal intervention dentistry. *International Dental Journal*, 68(3), 1–10.

- Selwitz, R. H., Ismail, A. I., & Pitts, N. B. (2019). Dental caries: Prevention and management strategies. *The Lancet Oral Health Series*, 394, 249–260.
- Kaur, S., Kaur, K., & Sharma, A. (2022). Role of digital dentistry in preventive oral healthcare. *Journal of Oral Biology and Craniofacial Research*, 12(4), 540–547.
- Estai, M., Kanagasingham, Y., Huang, B., & Bunt, S. (2020). The role of tele-dentistry and digital technologies in improving oral health access. *Journal of Telemedicine and Telecare*, 26(3), 147–155.
- Irfan, M., Delgado, R., & Chen, R. (2022). Smartphone-based dental imaging systems for oral health screening. *Digital Health*, 8, 1–12.
- Tonetti, M. S., Jepsen, S., Jin, L., & Otomo-Corgel, J. (2018). Impact of oral diseases on global health. *Journal of Clinical Periodontology*, 45(20), 1–8.
- Peres, M. A., Macpherson, L. M. D., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., & Watt, R. G. (2019). Oral diseases: A global public health challenge. *The Lancet*, 394(10194), 249–260.
- Tavares, M., & Lindefeld Caldeira, E. (2021). Preventive approaches for early childhood caries management. *Pediatric Dentistry*, 43(2), 115–123.
- Cvikl, B., Moritz, A., & Bekes, K. (2018). Diagnosis and management of early childhood caries. *Clinical Oral Investigations*, 22, 1773–1782.
- Gimenez, T., Braga, M. M., & Raggio, D. P. (2022). Advances in non-invasive caries detection technologies. *Caries Research*, 56(4), 321–333.
- Innes, N. P. T., Chu, C. H., Fontana, M., Lo, E. C. M., Thomson, W. M., & Uribe, S. (2020). A century of change towards prevention and minimally invasive dentistry. *Journal of Dental Research*, 99(6), 611–617.
- Featherstone, J. D. B., & Chaffee, B. W. (2022). The evidence for caries management by risk assessment and prevention strategies. *Advances in Dental Research*, 33(1), 3–10.

